

## UG20 Primary Anti-Reflux Surgery (Fundoplication)

Expires end of February 2023

If you need more information, please contact the department directly.

If you would like this information in different languages or formats (e.g. audio, Braille or large print), please ask a member of the healthcare team.

You can also contact:

---

---

---

You can get more information from [www.aboutmyhealth.org](http://www.aboutmyhealth.org)

Tell us how useful you found this document at [www.patientfeedback.org](http://www.patientfeedback.org)



Royal College  
of Surgeons  
of England



THE ROYAL  
COLLEGE OF  
SURGEONS  
OF EDINBURGH

**BBUGSS**

British Benign Upper  
Gastrointestinal Surgical Society



Patient Information Forum

[eidohealthcare.com](http://eidohealthcare.com)

**EIDO**  
HEALTHCARE

UNITED KINGDOM

## What is acid reflux?

Acid reflux is a condition where acid from your stomach passes up into your oesophagus (gullet). It is normal for a small amount of acid to travel into your oesophagus. If this happens too often it can cause symptoms of a burning sensation in your chest (heartburn) or acid in the back of your mouth. The acid can cause the lining of your oesophagus to become inflamed (oesophagitis) or scarred.

Your surgeon has suggested an operation to prevent the acid from travelling into your oesophagus. This document will give you information about the benefits and risks to help you to make an informed decision.

If you have any questions that this document does not answer, it is important that you ask your surgeon or the healthcare team.

Once all your questions have been answered and you feel ready to go ahead with the procedure, you will be asked to sign the informed consent form. This is the final step in the decision-making process. However, you can still change your mind at any point.

## How does acid reflux happen?


At the join between your stomach and oesophagus there is a weak valve that prevents acid and other stomach contents from travelling up into your oesophagus. Sometimes this valve

does not work effectively, causing acid reflux.


Your oesophagus normally passes through a natural gap in your diaphragm, this also contributes to the valve at the bottom of the oesophagus that prevents acid reflux. The majority of patients have the presence of a small hiatus hernia which can widen this gap and also weakens the strength of the valve which contributes to symptoms.

## What is a fundoplication?

A fundoplication procedure involves repairing any hiatus hernia that may be present as well as any excess gap in the diaphragm. Your surgeon will wrap the top part of your stomach around your oesophagus to create a valve to stop acid getting into the gullet. The operation is sometimes called a 'wrap'. There are different types of wrap, ranging from complete wrapping of the stomach around the oesophagus to partial wrapping. Your surgeon will talk to you about the most suitable procedure for you.



A normal valve



A faulty valve

## What are the benefits of surgery?

This surgery aims to improve your reflux symptoms. You should get relief from symptoms of acid reflux and no longer need to take medication.

## Are there any alternatives to surgery?

Avoiding foods that make your symptoms worse, especially late at

night can help ease your symptoms. Raising the head of the bed can also help. If you are overweight, it may help to lose weight.

Medication that lowers the acid content in your stomach is effective at controlling symptoms and healing the inflammation in your oesophagus. Medication called proton pump inhibitors is currently the most effective and is the main treatment for acid reflux.

There is a surgical alternative to a fundoplication called a LINX™ procedure. This involves placing magnetic beads around your oesophagus to form a valve effect. If you are suitable for this procedure, your surgeon will discuss it with you.

There are treatments available that use a telescope. These are not always available and long-term outcomes are less clear.

Surgery is only recommended if medication does not help, you have side effects from anti-acid medication or if you would prefer to have an operation than take medication for the rest of your life.

## What will happen if I decide not to have the operation?


Surgery is not essential and you can continue on the medication, particularly if your symptoms are well controlled.

It is important to follow the eating and drinking instructions that your doctor gives you and try to maintain a healthy weight. You should eat smaller meals more often and avoid chocolate, caffeine and alcohol. Try to eat at regular times and not in the 2 hours before you go to sleep.

## What does the operation involve?

The healthcare team will carry out a number of checks, including for any allergies and to make sure you have the operation you came in for. You can help by confirming to your surgeon and the healthcare team your name and the operation you are having.

The operation is performed under a general anaesthetic and usually takes 1 to 2 hours. You may also have injections of local anaesthetic to help with the discomfort after the operation. You may be given antibiotics during the operation to reduce the risk of infection.



A hiatus hernia

Your surgeon will hold your liver out of the way and free up the upper stomach and lower oesophagus, along with the muscular part of your diaphragm.

They will stitch your diaphragm to reduce the size of the hole your oesophagus passes through.

Your surgeon will wrap and stitch the top part of your stomach around your lower oesophagus, to produce a valve effect.


Your surgeon can wrap your stomach all the way round your oesophagus or just part-way round. Partial wraps can be performed behind or in front of your oesophagus. Your surgeon will tell you the best wrap to perform for you.

## Laparoscopic (keyhole) surgery

Your surgeon will use laparoscopic (keyhole) surgery as this is associated

with less pain, less scarring and a faster return to normal activities.

Your surgeon will make a cut on or near your belly button so they can insert an instrument in your abdominal cavity to inflate it with gas (carbon dioxide). They will make several small cuts on your abdomen so they can insert tubes (ports) into your abdomen.



Laparoscopic surgery

Your surgeon will hold your liver out of the way and free up the upper stomach and lower oesophagus, along with the muscular part of your diaphragm.

If you have a large gap in the diaphragm where the oesophagus passes through, your surgeon may stitch your diaphragm to reduce the large gap. This will improve your symptoms and reduce the chance of a hiatus hernia developing in the future.

Your surgeon will wrap and stitch the top part of your stomach around your

lower oesophagus, to reproduce a valve effect.

For less than 2 in 100 people it will not be possible to complete the operation using keyhole surgery. The operation may be changed (converted) to open surgery, which involves a larger cut on your upper abdomen and also results in a longer recovery period.

Your surgeon will remove the instruments and close the cuts.

### Open surgery

The operation is the same but it is performed through a larger cut on your upper abdomen. Sometimes your surgeon may recommend that the operation is performed through a cut on your chest.

Your surgeon may decide that keyhole surgery is not appropriate for you and recommend open surgery. They will discuss the reasons with you.

### What should I do about my medication?

Make sure your healthcare team knows about all the medication you take and follow their advice. This includes all blood-thinning medication as well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter.



## What can I do to help make the operation a success?

If you smoke, stopping smoking now may reduce your risk of developing complications and will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight. Your surgeon may suggest you follow a special diet for 2 weeks before the procedure to reduce the size of your liver. The liver is a large organ that needs to be lifted to perform the surgery safely. If it is smaller, the risk of complications such as bleeding are reduced.

Regular exercise should help to prepare you for the operation, help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

You can reduce your risk of infection in a surgical wound.

- In the week before the operation, do not shave or wax the area where a cut is likely to be made.
- Try to have a bath or shower either the day before or on the day of the operation.
- Keep warm around the time of the operation. Let the healthcare team know if you feel cold.

- If you are diabetic, keep your blood sugar levels under control around the time of your procedure.

If you have not had the coronavirus (Covid-19) vaccine, you may be at an increased risk of serious illness related to Covid-19 while you recover. Speak to your doctor or healthcare team if you would like to have the vaccine.

## What complications can happen?

The healthcare team will try to reduce the risk of complications.

Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you. Some risks are higher if you are older, obese, you are a smoker or have other health problems. These health problems include diabetes, heart disease or lung disease.

Some complications can be serious and can even cause death.

You should ask your doctor if there is anything you do not understand.

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

## General complications of any operation

- Bleeding during or after the operation. You may need a blood transfusion or another operation.
- Developing a hernia in the scar or port site. This appears as a bulge or rupture called an incisional hernia. If this causes problems, you may need another operation.
- Infection of the surgical site (wound). It is usually safe to shower after 2 days but you should check with the healthcare team. Let the healthcare team know if you get a high temperature, notice pus in your wound, or if your wound becomes red, sore or painful. An infection usually settles with antibiotics but you may need special dressings and your wound may take some time to heal. In some cases another operation might be needed. Do not take antibiotics unless you are told you need them.
- Allergic reaction to the equipment, materials or medication. The healthcare team is trained to detect and treat any reactions that might happen. Let your doctor know if you have any allergies or if you have reacted to any medication or tests in the past.
- Blood clot in your leg (deep-vein thrombosis – DVT). This can cause pain, swelling or redness in your leg, or the veins near the surface of your

leg to appear larger than normal. The healthcare team will assess your risk. They will encourage you to get out of bed soon after the operation and may give you injections, medication, or special stockings to wear. Let the healthcare team know straight away if you think you might have a DVT.

- Blood clot in your lung (pulmonary embolus), if a blood clot moves through your bloodstream to your lungs. Let the healthcare team know straight away if you become short of breath, feel pain in your chest or upper back, or if you cough up blood. If you are at home, call an ambulance or go immediately to your nearest Emergency department.
- Chest infection. If you have the operation within 6 weeks of catching Covid-19, your risk of a chest infection is increased (see the 'Covid-19' section for more information).
- Death (risk: less than 1 in 300).

## Specific complications of this operation

### Keyhole surgery complications

- Damage to structures such as your bowel, liver or blood vessels when inserting instruments into your abdomen (risk: less than 3 in 1,000). The risk is higher if you have had previous surgery to your abdomen. If an injury does happen, you may need open surgery. About 1 in 3 of these

injuries is not obvious until after the operation.

- Developing a hernia near one of the cuts used to insert the ports (risk: 1 in 100). Your surgeon will try to reduce this risk by using small ports (less than a centimetre in diameter) where possible or, if they need to use larger ports, using deeper stitching to close the cuts.
- Surgical emphysema (a crackling sensation in your skin caused by trapped carbon dioxide), which settles quickly and is not serious.
- Gas embolism. This is when gas (carbon dioxide) gets into the bloodstream and blocks a blood vessel. This is very rare but can be serious.

### Fundoplication complications

- Pneumothorax, where air escapes into the space around your lung. Sometimes the air will need to be let out by inserting a tube in your chest (chest drain).
- Making a hole in your oesophagus or stomach (perforation), which needs repairing (risk: 1 in 100). If this happens your healthcare team may leave a drain in to secure the repair. You may need to stay in hospital for longer to recover.
- Tear of the stitches used for the wrap, if you retch (strain to be sick) or vomit in the first few weeks. This may cause the wrap to become loose. Sometimes a tear can make a hole in

your stomach that will need to be repaired by surgery straight away.

- Damage to your liver when holding it out of the way (risk: 5 in 100). If the damage is serious, you may need another operation.
- Damage to your spleen. Your spleen may need to be removed.
- Difficulty swallowing for a few months because the site where your stomach is wrapped around your oesophagus is inflamed. This is common and you should be able to swallow most foods normally by 3 months.

### Long-term problems

- Continued difficulty swallowing where you cannot swallow most foods normally (risk: 5 in 100). If you find that food such as bread and meat get stuck, avoid them.
- Incomplete control of reflux symptoms, if the wrap is not tight enough or becomes loose (risk: less than 5 in 100). This may settle with medication.
- Weight loss during the first 2 months. It is normal to feel fuller than usual and you may be able to eat only small meals. Sit up when you eat and take a drink with your meal to help the food go down. Eat more often than before to try to keep your weight up. If you do lose weight, you will usually put it back on. If you have any



concerns about your diet, ask the dietician.

- Abdominal discomfort (risk: up to 3 to 5 in 10). You will probably not be able to burp as usual, which can cause gas to build up in your abdomen (gas bloat). You may pass more wind than usual.

- Diarrhoea (risk: less than 3 in 100). If loose or more frequent stools are troublesome, your doctor may give you some medication to slow down your bowel.

- Tissues can join together in an abnormal way (adhesions) when scar tissue develops inside your abdomen. Adhesions do not usually cause any serious problems but can lead to bowel obstruction. The risk is lower if you have keyhole surgery.

If any of these problems are severe and continue for over 3 months, you may need another operation (risk: less than 5 in 100). If you have these symptoms for over 3 months, let your surgeon know.

## Covid-19

A recent Covid-19 infection increases your risk of lung complications or death if you have an operation under general anaesthetic. This risk reduces the longer it is since the infection. After 7 weeks the risk is no higher than someone who has not had Covid-19. However, if you still have symptoms the risk remains high. The risk also depends on your age, overall

health and the type of surgery you are having.

You must follow instructions to self-isolate and take a Covid-19 test before your operation. If you have had Covid-19 up to 7 weeks before the operation you should discuss the risks and benefits of delaying it with your surgeon.

## Consequences of this procedure

- Pain. The healthcare team will give you medication to control the pain and it is important that you take it as you are told so you can move about and cough freely. After keyhole surgery, it is common to have some pain in your shoulders because a small amount of carbon dioxide gas may be left under your diaphragm. Your body will usually absorb the gas naturally over the next 24 hours, which will ease the symptoms.

- Scarring of your skin. In some cases this can be unsightly.

## How soon will I recover?

### In hospital

After the operation you will be transferred to the recovery area and then to the ward.

You will be given anti-sickness medication. You will be able to drink from the first day and then you will go on a soft diet. You should no longer need to take your acid-reducing medication.

You may be able to go home the same day or the next day. However, your doctor may recommend that you stay a little longer.

You need to be aware of the following symptoms as they may show that you have a serious complication.

- Pain that gets worse over time or is severe when you move, breathe or cough.
- A high temperature or fever.
- Dizziness, feeling faint or shortness of breath.
- Feeling sick or not having any appetite (and this gets worse after the first 1 to 2 days).
- Not opening your bowels and not passing wind.
- Swelling of your abdomen.
- Difficulty passing urine.

If you do not continue to improve over the first few days, or if you have any of these symptoms, let the healthcare team know straight away. If you are at home, contact your surgeon or GP. In an emergency, call an ambulance or go immediately to your nearest Emergency department.

## Returning to normal activities

If you had sedation or a general anaesthetic and you do go home the same day:

- a responsible adult should take you home in a car or taxi and stay with you for at least 24 hours;

- you should be near a telephone in case of an emergency;
- do not drive, operate machinery or do any potentially dangerous activities (this includes cooking) for at least 24 hours and not until you have fully recovered feeling, movement and co-ordination; and
- do not sign legal documents or drink alcohol for at least 24 hours.

To reduce the risk of a blood clot, make sure you carefully follow the instructions of the healthcare team if you have been given medication or need to wear special stockings.

You will need to eat slowly and chew your food thoroughly. Start with a liquid diet and build up to soft food. Your healthcare team will provide you with further guidance on what should be eating after your operation. You can normally return to a normal diet after 6 weeks. You will not be able to have fizzy drinks at all.

You should be able to return to work after a few weeks, depending on your recovery and the type of work you do.

Your doctor may tell you not to do any manual work for a while. Do not lift anything heavy for a few weeks.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Do not drive until you can control your vehicle, including in an

emergency, and always check your insurance policy and with the healthcare team.

Ask your healthcare team if you need to do a Covid-19 test when you get home.

## The future

You should make a full recovery, with the symptoms of acid reflux gone or much improved.

## Summary

Acid reflux can cause heartburn or acid in your mouth. The acid can cause the lining of your oesophagus to become inflamed or scarred. Surgery may be recommended if your symptoms continue while you are taking medication.

Surgery is usually safe and effective but complications can happen. You need to know about them to help you to make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you, and they may be able to tell you about any other suitable treatments options.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

## Acknowledgements

Reviewers: British Benign Upper Gastrointestinal Surgical Society, Ian Beckingham (DM, FRCS), Marcos Kostalas (MD FRCS), Stuart Andrews (MD FRCS), Simon Parsons (DM, FRCS)  
Illustrator: Medical Illustration Copyright © Medical-Artist.com